

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

## **IMAGES ARE BEST AVAILABLE COPY.**

As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.

**CAENORHABDITIS ELEGANS CHEMOSENSORY BIOASSAY FOR  
SEVEN TRANSMEMBRANE RECEPTOR LIGANDS**

**Cross-Reference to Related Application**

5           This application claims benefit to the earlier provisional U.S. Application No. 60/237,290, filed on October 2, 2000, the contents of which are incorporated herein by reference in their entirety.

**Field of the Invention**

10           This invention relates to methods for producing a transgenic *Caenorhabditis elegans* (*C. elegans*) that expresses a human 7-transmembrane receptor (herein "7TMR") in its neurons that correlate with behavior. These transgenic *C. elegans* nematodes can be used in a variety of ways, including, but not limited to: (1) screening and identifying substances that bind to and activate human 7TMRs; (2) screening for substances that  
15           antagonize human 7TMRs; (3) identifying 7TMRs that may respond to particular substances; and (4) evaluating the specificity and efficacy of substances on human 7TMR activation.

**Background of the Invention**

20           With the increasing flow of DNA sequence information from a variety of species, including humans, there is a need for new methods that can link function with gene sequence. We have developed an invention useful for probing the function of cell surface, transmembrane receptors, including the largest, and most pharmacologically important, gene classes in multicellular organisms, the seven transmembrane receptor (herein referred  
25           to as "7TMR") class of genes. *See generally* Stadel, *et al.*, *Trends Pharmacol. Sci.* 18(11): 430-7 (1997). Receptors often reside at the membrane surface of the cell and, upon binding of a ligand, activate an intracellular signaling cascade that mediates a particular cellular response. A crucial step in understanding the function of a particular receptor is to identify the ligand, or a surrogate ligand, that can bind and activate the receptor. The invention we  
30           describe here, which we dub the "*C. elegans* Chemosensory Bioassay," allows nematodes to sense and respond to substances that may affect cell surface receptor activity. This invention can be used in a variety of ways, including, but not limited to: (1) screening and identifying substances that bind to and activate human 7TMRs; (2) screening for substances that antagonize human 7TMR activation; (3) identifying human 7TMRs that may respond to

particular substances; and (4) evaluating the specificity and efficacy of substances on human 7TMR activation.

The instant invention represents a significant improvement over current technologies for probing transmembrane receptor function. Current methods for identifying activating substances for receptors is, at best, medium throughput (*See Stadel, et al., supra*). Elaborate and sophisticated molecular assays designed to measure such parameters as calcium mobilization, cAMP, GTP- $\gamma$  S binding, inositol phosphate production, MAP kinase activation, *etc.*, are used to identify activating substances. These assays involve either sophisticated machines that can detect receptor activation in transgenic mammalian cell lines (Sullivan, *et al., Methods Mol Biol* 114: 125-33 (1999)) or labor-intensive methods, such as microinjection of *Xenopus* oocytes followed by electrophysiological recording (Wagner, *et al., Cell Physiol Biochem.* 10(1-2): 1-12 (2000)) Such assays involve direct mechanical detection of receptor function.

By contrast, our invention links receptor activation to simple nematode behaviors, obviating the need for sophisticated instrumentation. In addition, the self-replicating biological nature of nematodes allows large numbers of animals to be produced in the laboratory, ensuring that our invention will be an inexpensive solution and will be amenable to high-throughput applications. For example, libraries of substances can be easily screened, as discussed below. Also, unlike mechanical detection of receptor activation, our invention does not require any prior assumptions about the parameters to be measured (calcium, cAMP, *etc.*). Instead, our invention uses simple biological behaviors in nematodes to assay receptor activation.

Cell surface receptors are a large collection of proteins that fit into a number of superfamilies by their structural similarities. These superfamilies include, but are not limited to: ion channels, transporters, and 7 transmembrane receptors (Saier MH Jr, *J Membr Biol* 175(3):165-180 (2000); Saier MH Jr, *J Cell Biochem, Suppl* 32-33:84-94 (1999); Bockaert J, Pin JP, *EMBO* 18(7):1723-1729 (1999)). It is well established that many medically significant biological processes are mediated by such proteins. For example, the most common inherited disease among caucasians is cystic fibrosis, caused by mutations in the CFTR chloride channel (Frizzell RA *Physiol Rev*, 79(1 Suppl):S1-2 (1999)). In addition, a number of other disease states are linked to mutations in other ion channels (Weinreich F, Jentsch TJ, *Curr Opin Neurobiol* 10(3):409-415 (2000)). One role of the transporter class of cell surface proteins is the uptake of neurotransmitters from synaptic clefts in the nervous system. These transporters are targets for therapeutic

intervention in a variety of psychiatric disorders. For example, selective serotonin uptake inhibitors are used in the treatment of a number of disorders and include the widely prescribed drug Prozac (Masand PS, Gupta S, *Harv Rev Psychiatry*, 7(2):69-84 (1999)).

Among the cell surface receptor superfamilies, the 7 transmembrane receptor (7TMR) superfamily appears to be the largest in metazoan organisms (*see* Bockaert and Pin, *supra*). 7TMRs mediate a variety of cellular responses to extracellular stimuli. For example, 7TMRs bind or interact with a huge variety of ligands; including photons, small molecules, and large proteins (*see* Bockaert and Pin, *supra*). 7TMRs are be intracellularly coupled by heterotrimeric G-proteins to various intracellular enzymes, ion channels and transporters (*see*, Johnson, *et al.*, *Endoc. Rev.*, 1989, 10:317-331). The 7TMRs include dopamine receptors, which bind to neuroleptic drugs used for treating psychotic and neurological disorders. Other examples of members of this family include, but are not limited to, calcitonin, adrenergic, endothelin, cAMP, adenosine, muscarinic, acetylcholine, serotonin, histamine, thrombin, kinin, follicle stimulating hormone, opsins, endothelial differentiation gene-1, rhodopsins, odorant, and cytomegalovirus receptors. Thus, these receptors mediate a large variety of biological and cellular responses to extracellular cues and include a wide range of biologically active receptors, such as hormone, viral, growth factor and neuroreceptors.

For most cell surface receptors identified in genome sequencing efforts, the activating ligand is not known. However, given the wide variety and important nature of biological processes regulated by cell surface receptors, determination of the activating ligand is of fundamental pharmaceutical importance. For example, over the past 15 years nearly 350 therapeutic agents targeting 7TMRs have been successfully introduced onto the market (*See* Stadel, *et al.*, *supra*). Therefore, finding out what ligands bind these 7TMRs will enable researchers to develop drugs to either agonize or antagonize the interaction of 7TMRs with their ligands. Our invention can be applied to this problem and is useful for probing the function of a large number of cell surface protein types by linking simple *C. elegans* behaviors to the activation of human 7TMRs.

### Summary of the Invention

In one aspect, the invention relates to a method for producing a transgenic *Caenorhabditis elegans* (*C. elegans*) that expresses at least one human 7TMR pan-neuronally, such that said transgenic *C. elegans* exhibits a known phenotype, said method comprising the steps of:

(a) producing a transgene by operably linking a gene expression construct that encodes a human 7TMR to a pan-neuronal promotor; and

(b) introducing said transgene into said transgenic *C. elegans*, such that said transgenic *C. elegans* expresses said human 7TMR pan-neuronally; and

5 (c) examining said *C. elegans* to determine that said *C. elegans* exhibits a known phenotype.

During the practice of this method, wherein the known phenotype is selected from the group consisting of: exploded (Exp), dumpy (Dpy), long body (Lon), hyperactive movement (Hpr), paralyzed (Prl), molt defect (Mlt), sterile (Ste), sick (Sck), body  
10 morphology defect (Bmd), vulvaless (Vul), slow growth (Gro), egg laying defect (Egl), larval arrest (Lva), larval lethal (Let), protruding vulva (Pvl), multiple vulva (Muv), sterile progeny (Stp), small (Sma), clear (Clr), blistered (Bli), high incidence of male progeny (Him), roller (Rol), larval lethal (Lvl), uncoordinated (Unc), embryonic lethal (Emb).

In another aspect, this invention provides a transgenic *C. elegans* produced by the above  
15 method.

In one aspect, the invention relates to a method for producing a transgenic *Caenorhabditis elegans* that expresses a human 7-transmembrane receptor (7TMR) in the sensory neurons that correlate with behavior, said method comprising the steps of:

(a) producing a transgene by operably linking a gene expression construct that  
20 encodes a human 7TMR to a sensory neuron promotor; and

(b) introducing said transgene into said *C. elegans*, such that said *C. elegans* expresses said human 7TMR in its sensory neurons that correlate with behavior.

The gene expression construct used in this method can be either DNA or a plasmid. The behavior referred to in this method is selected from the group consisting of: water-soluble  
25 chemorepulsion, water-soluble chemoattraction, volatile chemorepulsion, volatile chemoattraction, thermorepulsion, thermoattraction, and dauer formation.

In one aspect of this method, the behavior is water-soluble chemorepulsion, and said sensory neurons are selected from the group consisting of: ASH and ADL neurons. In this method, the sensory neuron promotor is the promotor region of *gpa-11*, and the plasmid  
30 construct is pCEASH2.1.

In another aspect, the instant invention relates to the above-referenced method, wherein said behavior is water-soluble chemoattraction, and said sensory neurons are selected from the group consisting of: ASE, ADF, ASG, and ASI neurons. In this method, the sensory neuron promotor is the promotor region of a gene selected from the group consisting of: *gcy-5*, *gcy-6*, and *gcy-7*.

In yet another aspect of this method, the behavior is volatile chemorepulsion, and said sensory neurons are AWB neurons. In this embodiment of the invention, the sensory neuron promotor is the promotor region of *str-1*.

In another aspect of this method, the behavior is volatile chemoattraction, and the sensory neurons are AWA neurons. In this embodiment, the sensory neuron promotor is the promotor region of *odr-10*.

In another aspect of this method, the behavior is dauer formation, and said sensory neurons are selected from the group consisting of: ASI, ASG, and ADF neurons.

In yet another aspect of this method, the behavior is selected from the group consisting of: thermoattraction and thermorepulsion, and said sensory neurons are AFD neurons. In this embodiment, the sensory neuron promoter is the promoter region of *gcy-8*.

In another aspect, this method of producing transgenic *C. elegans* further comprises the step of:

(c) introducing a DNA construct that expresses a reporter gene with a nematode promotor.

Preferably, this reporter gene encodes a fluorescent protein, such as a green fluorescent protein (GFP).

In another aspect, this invention provide a method for producing a transgenic *C. elegans* that expresses a human 7TMR receptor in the sensory neurons that correlate with behavior, said method comprising the steps of:

(a) producing a first transgene by operably linking a gene expression construct that encodes a human 7TMR to a sensory neuron promotor;

(b) producing a second transgene by comprising an accessory protein operably linked to a promotor; and

(c) introducing said first and second transgenes into said *C. elegans*, such that said *C. elegans* coexpresses both said accessory protein and said human 7TMR in its sensory neurons that correlate with behavior.

In one embodiment of this method, the non-nematode cell surface receptor is a 7TMR, and said accessory protein is a human G protein.

Another embodiment of this invention encompasses a transgenic *C. elegans* produced by any of the above-described methods.

In yet another aspect, this invention provides a method for identifying at least one ligand of at least one human 7TMR, said method comprising the steps of:

- 5 (a) providing at least one *C. elegans* that expresses at least one human 7TMR in the sensory neurons that correlate with behavior;
- (b) contacting said at least one *C. elegans* with at least one test substance; and
- (c) detecting modulation of behavior of said at least one *C. elegans* in response to said at least one test substance.

- 10 In this screening method, the behavior is selected from the group consisting of: water-soluble chemorepulsion, water-soluble chemoattraction, volatile chemorepulsion, volatile chemoattraction, thermorepulsion, thermoattraction, and dauer formation.

In another aspect, the instant invention provides a method for identifying at least one ligand of at least one human 7TMR, said method comprising the steps of:

- 15 (a) providing at least one *C. elegans* that expresses at least one 7TMR in its sensory neurons that correlate with water-soluble chemorepulsive behavior;
- (b) placing a medium into a first portion of a receptacle, wherein said receptacle is divided into at least two portions;
- (c) adding at least two different concentrations of at least one chemorepulsant
- 20 substance to the first portion of said receptacle;
- (d) adding said at least one *C. elegans* to the second portion of said receptacle; and
- (e) detecting, after a suitable time period, the behavioral response of said at least one *C. elegans* to said at least one chemorepulsant substance.

- In yet another aspect, this screening method further comprises the following step:
- 25 (f) generating a chemotactic index, wherein said chemotactic index is the fraction of said at least one *C. elegans* that moved to said first portion of said receptacle.

In another aspect of this method, said sensory neurons are selected from the group consisting of: ASE, ADF, ASG and ASI neurons.

- In this method, said at least one chemorepulsive substance is selected from the group
- 30 consisting of: copper sulfate, sodium dodecyl sulfate (SDS), D-tryptophan, heptanol, octanol, nonanol, nonanone, benzaldehyde, 2, 4, 5-trimethylthiazole, and ethyl heptanoate.

In one embodiment of this screening method, the growth medium comprises a biomolecular separation in a matrix. This matrix is selected from the group consisting of: agarose and polyacrylamide.

In one aspect of this screening method, said at least one test substance can be present in beads on a solid or semi-solid medium.

In other aspects of this screening method, said at least one test substance comprises at least one natural peptide, a viral plaque, or is secreted by at least one microorganismal colony. In another aspect, said at least one test substance is a recombinant peptide that is secreted by at least one microorganismal colony. Said at least one recombinant peptide can be a peptide library. Said microorganismal colony is selected from the group consisting of: bacteria and yeast.

In yet another aspect of this screening method, said detecting step (e) is carried out by direct detection with a suitable optical instrument. Said detecting step (e) can be carried out by assisted detection of the optical density over the surface of said medium. Alternatively, said assisted detection can be mechanical detection.

In a yet another aspect, the instant invention provides a method for identifying at least one ligand of at least one human 7TMR, said method comprising the steps of:

- (a) providing at least one *C. elegans* that expresses at least one human 7TMR in its sensory neurons that mediate water-soluble chemoattractive behavior;
- (b) placing a medium into a first portion of a receptacle, wherein said receptacle is divided into at least two portions;
- (c) adding at least two different concentrations of at least one chemoattractant substance to the first portion of said receptacle;
- (d) adding said at least one *C. elegans* to the second portion of said receptacle; and
- (e) detecting, after a suitable time period, the behavioral response of said at least one *C. elegans* to said at least one chemoattractant substance.

In another aspect of this method, said sensory neurons are selected from the group consisting of: ASE, ADF, ASG and ASI neurons.

In yet another aspect of this invention, this method further comprises the following step:

- (f) generating a chemotactic index, wherein said chemotactic index is the fraction of said at least one *C. elegans* that moved to said first portion of said receptacle.

In one embodiment of the instant invention, said at least one chemoattractant substance is selected from the group consisting of: isoamyl alcohol, biotin, lysine, histidine, cysteine, and serotonin.

In another aspect, this invention provides a method for identifying at least one ligand of a human 7TMR said method comprising the steps of:



(a) providing at least one *C. elegans* that expresses a human 7TMR in the sensory neurons that correlate with behavior;

(b) placing at least one test substance on a substrate surface that contains growth medium;

5 (c) placing a uniform lawn of bacteria onto the surface of said growth medium;

(d) contacting said at least one *C. elegans* with the surface of said growth medium; and

(e) detecting, after a suitable time period, a decrease in the density of said uniform lawn of bacteria.

10 The bacteria employed in this method is selected from the group consisting of:

*Acinetobacter calcoaceticus*, *Bacillus cereus*, *Bacillus sp.*, *Enterobacter amnigenus*,  
*Enterobacter cloacae*, *Escherichia coli*, *Flavobacterium sp.*, *Proteus vulgaris*,  
*Pseudomonas aeruginosa*, *Pseudomonas cepacia*, *Pseudomonas maltophilia*, *Pseudomonas putida*, *Pseudomonas sp.*, *Serratia liquefaciens*, *Serratia marcesens*, *Zymobacillus*  
 15 *macerans*. Preferably, the bacteria used in this method is *E. coli*.

In yet another aspect of this method, the detection step (e) comprises photometrically determining variations in the density of said bacterial lawn.

In a further aspect of this screening method, the bacteria express GFP, and the detecting step (e) comprises determining the fluorescence intensity of said bacterial lawn.

20 In yet another aspect, the instant invention provides a method for identifying at least one ligand of at least one human 7TMR, said method comprising the steps of:

(a) providing at least one *C. elegans* that expresses a human 7TMR in the sensory neurons that correlate with behavior;

(b) placing a medium in a receptacle;

25 (c) arranging at least one test substance in said receptacle;

(d) adding said at least one *C. elegans* to said receptacle; and

(e) detecting, after a suitable time period, the behavioral response of said at least one *C. elegans* over the surface of said medium.

30 In another aspect, the instant invention provides a method for evaluating the potency of human 7TMR activation by a known ligand, said method comprising the steps of:

(a) providing at least one *C. elegans* that expresses at least one human 7TMR in the sensory neurons that correlate with behavior, wherein said at least one human 7TMR has a known ligand;

(b) contacting said at least one *C. elegans* with said ligand and at least one structurally related compound; and

(c) detecting the behavioral response of said at least one *C. elegans* to said at least one structurally related compound; and

5 (d) comparing the behavioral response of said at least one *C. elegans* to said ligand to the behavioral response of said at least one *C. elegans* to said at least one structurally related compound.

In one aspect of this method, the behavior is selected from the group consisting of: chemoattraction and chemorepulsion.

10 In another aspect, this invention relates to a method for identifying at least one test substance that modulates the activation of at least one human 7TMR by a known ligand, said method comprising the steps of:

(a) providing at least one *C. elegans* that expresses at least one human 7TMR in the sensory neurons that correlate with behavior, wherein said at least one human 7TMR has a  
15 known ligand;

(b) contacting said at least one *C. elegans* with at least one test substance in the presence of said known ligand; and

(c) determining whether said at least one test compound binds to and activates or inhibits the activation of said at least one human 7TMR by said ligand by measuring the  
20 alteration of behavior of said at least one *C. elegans* in response to said at least one test substance.

In a further aspect of this method, the behavior that is measured is selected from the group consisting of: water-soluble chemorepulsion, water-soluble chemoattraction, volatile chemorepulsion, volatile chemoattraction, thermorepulsion, thermoattraction, and dauer  
25 formation.

In yet another aspect, this invention provides a method for identifying at least one test substance that is an antagonist of a human 7TMR, said method comprising the steps of:

(a) providing at least one *C. elegans* that expresses a human 7TMR pan-neuronally, wherein said human 7TMR is activated by an endogenous ligand, such that said transgenic  
30 *C. elegans* exhibits a known phenotype;

(b) contacting said at least one *C. elegans* with at least one test substance, wherein said at least one test substance is distributed in a medium;

(c) determining whether said at least one test substance causes a suppression of said known phenotype in said at least one transgenic *C. elegans*; and

(d) identifying said at least one test substance that causes a suppression of said known phenotype in said at least one transgenic *C. elegans* as an antagonist of said human 7TMR.

In another aspect, this invention provides a method for identifying a surrogate ligand present in a strain of transgenic *C. elegans* that expresses a human 7TMR pan-neuronally, wherein said strain of transgenic *C. elegans* exhibits a known phenotype, such that said human 7TMR is activated by an endogenous ligand, said method comprising the steps of:

(a) providing a strain of human 7TMR-expressing transgenic *C. elegans*, wherein said strain of transgenic *C. elegans* exhibits a known phenotype;

(b) subjecting said strain of transgenic *C. elegans* to at least one mutagenic screen; and

(c) determining whether said at least one mutagenic screen results in a suppression of said known phenotype in said strain of transgenic *C. elegans*.

In yet another aspect, this invention provides a method for identifying at least one substance that agonize the activity of a human 7TMR, said method comprising the steps of:

(a) providing at least one transgenic *C. elegans* that expresses a human 7TMR pan-neuronally, wherein said at least one transgenic *C. elegans* does not exhibit a known phenotype because said human 7TMR is not activated by an endogenous ligand;

(b) contacting said at least one transgenic *C. elegans* with at least one test substance, wherein said at least one test substance is distributed in a medium;

(c) determining whether said at least one test substance causes said at least one said transgenic *C. elegans* to exhibit a known phenotype; and

(d) identifying said at least one test substance that causes said at least one said transgenic *C. elegans* to exhibit a known phenotype as an agonist of said human 7TMR.

### Brief Description of the Drawings

Figure 1 diagrams the pCEASH2.1 plasmid, which is capable of driving expression in the ADL and ASH neurons of *C. elegans* that mediate water-soluble chemorepulsion. In brief, the promoter region of the *gpa-11* gene has been placed in a plasmid context with the 3' region of the *unc-54* gene (see example 1 text for a detailed description). This *gpa-11* promoter region is sufficient to drive expression in the *C. elegans* ADL and ASH neurons.

Figure 2 demonstrates a nematode chemotaxis assay using an array of test substances. A round petri plate has been filled with an solid agar substrate. Test substances were spotted as

indicated in panel B. A test population of *C. elegans* was placed in the center of the plate (see leftmost panels). The distribution of animals was determined at the initial time, and after 2 hours by visual inspection of the plate. The plate was also recorded by digital photography at the initial and 2 hour timepoints for this figure. In panel A, the actual digital photomicrographs are displayed. In panel B, the positions of the animals and the test substances is schematized based on the digital photomicrographs. The test population shows a clear preference for the spot of isoamyl alcohol.

Figure 3 demonstrates consumption of GFP-expressing bacteria by a culture of *C. elegans*. Two cultures of *C. elegans* and GFP-expressing bacteria were placed in microtiter wells. In one well, a lethal concentration of 10 nanomolar ivermectin was placed. The GFP fluorescence in the wells was determined by a fluorescent plate reader instrument at the initial time, and after two days of culture. Fluorescence is displayed as Relative Intensity normalized to the fluorescent intensity present in the initial ivermectin-containing well. Notably, in the ivermectin-containing well, in which the *C. elegans* animals are dead, GFP fluorescence increases as the bacteria grow. In contrast, the non-ivermectin well shows a decrease in GFP fluorescence as the *C. elegans* population consumes the bacteria.

Figure 4 diagrams the pCEPN plasmid, which is capable of driving expression in the nervous system of *C. elegans*, as described in Example 4. In brief, the promoter region of the unc-119 gene has been placed in a plasmid context with the 3' region of the unc-54 gene (see example 4 text for a detailed description). This unc-119 promoter region is sufficient to drive expression in the *C. elegans* nervous system.

Figure 5 illustrates Egl, or egg-laying, phenotypes produced by expression of human 7TMRs in the nervous system of *C. elegans* (see example 5 for a detailed description of receptors). In brief, 5-8 single transgenic animals were allowed to lay eggs for 40 hours. The number of eggs laied were counted and graphed. The N2 strain is the canonical wildtype non-transgenic *C. elegans* strain. The receptor-expressing strains are indicated below each bar by the name of the expressed receptor. In comparison to N2, most of the transgenic strains show a reduction in the number of eggs laid.

## Glossary

The following illustrative explanations are provided to facilitate understanding of certain terms used frequently herein. The explanations are provided as a convenience and are not limiting of the invention.

5           The term, "behavior," as used herein, refers to behavior selected from the group consisting of: volatile chemotaxis, water- soluble chemotaxis, thermotaxis, and dauer formation. More specifically, these behaviors encompass water-soluble chemorepulsion, water-soluble chemoattraction, volatile chemorepulsion, volatile chemoattraction, thermorepulsion, and thermoattraction.

10           The term, "chemoattractant substance," as used herein, refers to both known chemoattractants, such as isoamyl alcohol, biotin, lysine, histidine, cysteine, and serotonin (Bargmann CI., *Chemotaxis and Thermotaxis*, In Riddle, Blumenthal, Meyer, and Priess (eds.), *C. ELEGANS* II, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY (1997)), as well as substances that are discovered to elicit chemoattractive behavior from  
15 transgenic *C. elegans* of the instant invention.

          The term, "chemorepellant substance," as used herein, refers to both known chemorepellants, such as copper sulfate, sodium dodecyl sulfate (SDS), D-tryptophan, heptanol, octanol, nonanol, nonanone, benzaldehyde, 2, 4, 5-trimethylthiazole, and ethyl heptanoate (Bargmann, *supra*), as well as substances that are discovered to elicit  
20 chemorepulsive behavior from transgenic nematodes of the instant invention.

          As used herein, phrase "known phenotype," includes, but is not limited to, the following phenotypes that have been scored in *C. elegans*: exploded (Exp), dumpy (Dpy), long body (Lon), hyperactive movement (Hpr), paralyzed (Prl), molt defect (Mlt), sterile (Ste), sick (Sck), body morphology defect (Bmd), vulvaless (Vul), slow growth (Gro), egg  
25 laying defect (Egl), larval arrest (Lva), larval lethal (Let), protruding vulva (Pvl), multiple vulva (Muv), sterile progeny (Stp), small (Sma), clear (Clr), blistered (Bli), high incidence of male progeny (Him), roller (Rol), larval lethal (Lvl), uncoordinated (Unc), embryonic lethal (Emb).

          As used herein, the term, "operably linked," refers to a DNA sequence and a  
30 regulatory sequence(s) that are connected in such a way as to permit gene expression when the appropriate molecules, *e.g.*, transcriptional activator proteins, are bound to the regulatory sequence(s). A nucleic acid is "operably linked" when it is placed into a functional relationship with another nucleic acid sequence. For example, DNA for a

presequence or a secretory leader is operably linked to DNA for a polypeptide if it is expressed as a preprotein that participates in the secretion of the polypeptide; a promoter or enhancer is operably linked to a coding sequence if it affects the transcription of the sequence; or a ribosome binding site is operably linked to a coding sequence if it is positioned so as to facilitate translation. Generally, "operably linked" means that the DNA sequences being linked are contiguous and, in the case of a secretory leader, contiguous and in reading phase. However, enhancers do not have to be contiguous. Linking is accomplished by ligation at convenient restriction sites. If such sites do not exist, then synthetic oligonucleotide adaptors or linkers are used in accord with conventional practice.

As used herein, the term, "pan-neuronal" or "pan-neuronally" refers to the expression of a given gene in all of the neurons of a transgenic *C. elegans*.

The term, "structurally related compounds," as it is used herein, refers to compounds that share a common pharmacophore, which is a core structure of the compound that has biological activity. Such "structurally related compounds" can differ from each other, however, in their substituent groups. As used in the screening assays herein, "structurally related compounds" share a pharmacophore with a known or newly discovered ligand of a non-nematode cell surface receptor that is expressed in the transgenic nematodes described below.

The term, "test population," as it is used herein, refers to a group *C. elegans* from the same strain that each express the same human 7TMR(s).

The term, "test substance," as used herein, refers to an organic or inorganic molecule, a homogenous mixture of such molecules, biological extracts, or biological fractions. A "test substance" could be, for example, a viral plaque or a natural peptide. A "test substance" could be, for example, a known chemoattractant or chemorepellent compound or a molecule that is discovered to elicit chemosensory behavior from transgenic nematodes of the instant invention.

"Transformation" means introducing DNA into an organism so that the DNA is replicable, either as an extrachromosomal element or by chromosomal integration. Depending on the host cell used, transformation is done using standard techniques appropriate to such cells. The calcium treatment employing calcium chloride, as described by Cohen, S. N., *Proc. Natl. Acad. Sci. (USA)*, 69: 2110 (1972) and Mandel, *et al.*, *J. Mol. Biol.* 53: 154 (1970), is generally used for prokaryotes or other cells that contain substantial cell-wall barriers. For mammalian cells without such cell walls, the calcium phosphate

precipitation method of Graham, *et al.*, *Virology* 52: 456-457 (1978) is preferred. General aspects of mammalian cell host system transformations have been described by Axel in U.S. Pat. No. 4,399,216, issued Aug. 16, 1983. Transformations into yeast are typically carried out according to the method of Van Solingen, P., *et al.*, *J. Bacteriology* 130: 946 (1977) and Hsiao, C. L., *et al. Proc. Natl. Acad. Sci (USA)* 76: 3829 (1979). However, other methods for introducing DNA into cells such as by nuclear injection, electroporation or by protoplast fusion may also be used. Transformations into *C. elegans* can be carried out by microinjection, as described in Mello CC, *et al*, *EMBO J.* 10:3959-3970 (1991), or by microparticle bombardment: as described in Wilm T, *et al.*, *Gene* 229:31-35 (1999) and Praitis *et al.*, *Genetics* 157, 1217-26 (2001).

### Detailed Description of the Invention

In a particularly preferred embodiment of the invention, the roundworm *Caenorhabditis elegans* is employed. *C. elegans* is a simple soil nematode species that has been extensively described at the cellular and molecular level, and is a model organism for biological studies (See Riddle, *C. ELEGANS II.*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY. (1997); The *C. elegans* Sequencing Consortium, *Science* 282: 2012-2018. (1998); Sulston, WB (ed), *THE NEMATODE CAENORHABDITIS ELEGANS*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, pp. 123-155 (1988)). The adult hermaphrodite is composed of 959 cells, 302 of which are defined as neuronal cells. Sulston, *supra*. *C. elegans* can respond to a variety of substances in its environment by moving away from a point source, termed chemorepulsion, or moving toward a point source, termed chemoattraction. Bargmann, *supra*. *C. elegans* also responds to a pheromone in the environment by taking an alternative developmental pathway, the dauer stage. Riddle, *et al.*, *Genetic and Environmental Regulation of Dauer Larva Development*, In Riddle, Blumenthal, Meyer, and Priess (eds), *C. ELEGANS II.*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY. (1997). Much of this chemosensory behavior is mediated by sensory organs termed the amphids, a pair of bilaterally symmetric structures located at the anterior of the animal. Each amphid contains the endings of 12 sensory neurons, eight of which are directly exposed to the environment through an opening in the cuticle of the animal. Bargmann, *supra*. The cell bodies of the sensory neurons lie in the nerve ring of the animal, where they synapse with other neurons to pass on sensory information. The function of each amphid sensory neuron has been defined by laser ablation microsurgery and subsequent testing of the operated animal. *Id.* Each type of

sensory modulates a particular chemosensory behavior. For example, the AWA neuron mediates chemotaxis toward volatile compounds, including diacetyl, pyrazine, and thiazole; whereas the AWB neuron mediates chemorepulsive behavior toward volatile compounds. The ASI, ASG, and ADF neurons are necessary in the process of dauer pathway commitment. Riddle, *et al.*, *supra*. The AFD neurons mediate thermotactic behavior, which includes thermoattraction and thermorepulsion. Bargmann, *supra*. The observation that particular neurons are needed for different behaviors suggests that behavioral modalities are determined by the properties and synaptic connections of these neurons.

While chemotactic behavior is determined at the neuronal level, the ability to discriminate among substances in the environment is determined by expression of particular receptors in the chemosensory neurons. For example, a *C. elegans* 7TM receptor, *odr-10*, has been defined as a requisite receptor for the chemotactic response of *C. elegans* to diacetyl. Sengupta, *et al.*, *Cell* 84: 899-909 (1996). Animals that are mutant for the *odr-10* gene no longer respond to diacetyl, and expression of *odr-10* in mammalian tissue culture cells confers Ca<sup>2+</sup> mobilization in response to diacetyl, indicative of activation. Sengupta, *et al.*, *supra*; Zhang, *et al.*, *PNAS, USA* 94: 12162-67 (1997). In *C. elegans*, ODR-10 is normally expressed in the AWA neuron, which mediates chemoattraction to diacetyl. These results strongly indicate that the ligand for *odr-10* is diacetyl, and ligand binding and *odr-10* activation in the AWA neuron leads to a chemotactic response to diacetyl. *Odr-10* belongs to a large family of 7TM receptors encoded in the genome of *C. elegans*, and it is likely that expression of these receptors in the sensory neurons provides the wide range of chemotactic chemical sensitivity seen in *C. elegans*. Given that different neurons are involved in different chemotactic behaviors, the expression of a receptor in certain neurons likely determines the behavioral response to the substance recognized by the receptor. Researchers tested this hypothesis by misexpressing *odr-10* in the AWB neuron, which mediates volatile chemorepulsive behavior. Troemel, *et al.*, *Cell* 91:161-169 (1997). This misexpression of *odr-10* led to a "reprogramming" of the chemotactic response, such that the animals showed chemorepulsive behavior to diacetyl, instead of chemoattraction. These observations lead us to believe that ligand specificity is provided by the receptor itself, while the behavioral response is determined by which neuron the receptor is expressed in.

The ability to reprogram the chemosensory response of *C. elegans* is not limited to misexpressing *C. elegans* receptors. At a recent *C. elegans* worm meeting (Tobin, *et al.*, West Coast Worm Meeting, June 23-26, 2000; <http://elegans.swmed.edu/WCWM/2000/>), a second example of reprogramming the chemosensory response of *C. elegans* was presented



using a mammalian-derived receptor. Expression of the mammalian capsaicin receptor VR1 in sensory neurons of *C. elegans* conferred a chemoavoidance behavior in response to capsaicin. Like the *odr-10* example discussed above, this report is a demonstration that nematode chemosensory behavior in response to substances can be modified by expression of a receptor in the sensory neurons of nematodes. Unlike our invention, the authors do not express the human capsaicin receptor in *C. elegans* with the purpose of using chemotactic behavior as a way of identifying substances that activate the receptor, nor do they propose that such an application is possible. We propose that the modification of chemosensory behavior by human 7TMR expression in sensory neurons can be applied to the identification and characterization of substances that activate human 7TMRs or modify human 7TMR activity.

#### **Production of Transgenic *C. elegans***

The transgenic *C. elegans* of this invention can be used in a variety of ways, including, but not limited to: (1) screening and identifying substances that bind to and activate particular human 7TMRs; (2) screening for substances that antagonize human 7TMR activation; (3) identifying human 7TMRs that may respond to particular substances; and (4) evaluating the specificity and efficacy of substances on human 7TMR activation.

One preferred embodiment of the invention includes a method for producing a transgenic *C. elegans* that expresses a human 7TMR in the sensory neurons that correlate with behavior, said method comprising the steps of:

- (a) producing a transgene by operably linking a gene expression construct that encodes a human 7TMR to a sensory neuron promoter; and
- (b) introducing said transgene into said *C. elegans*, such that said *C. elegans* expresses said human 7TMR in the sensory neurons that correlate with behavior.

In one particularly preferred embodiment of the instant invention, transgenic *C. elegans* are used as biosensors for substances that activate human 7TMRs or that block human 7TMR activation. This is accomplished by specific expression of human 7TMR(s) in the appropriate sensory neurons of *C. elegans*, allowing the organisms to sense and respond in a chemosensory fashion to substances that activate the human 7TMR(s). By expressing human 7TMR(s) in particular sensory neurons, the response of the animals to receptor activation will be manifested as one of several behaviors: chemorepulsion, chemoattraction, thermotaxis, including thermoattraction and thermorepulsion, or commitment to the dauer developmental pathway. Chemorepulsion can be toward volatile

compounds, mediated by the AWB sensory neuron of *C. elegans*, or toward water-soluble compounds, mediated by the ASH and ADL sensory neurons. A human 7TMR can be expressed in the AWB neuron of *C. elegans* using the promoter region of the *str-1* gene operably linked to the receptor coding sequence (Troemel ER, *et al*, *Cell* 91:161-169 (1997)). A receptor can be expressed in the ASH and ADL neurons using the promoter region of the *gpa-11* gene operably linked to the receptor coding sequence (Jansen G, *et al*, *Nature Genetics* 21:414-419 (1999)). Chemoattraction can likewise be toward volatile compounds, mediated by the AWA neuron, or toward water-soluble compounds, mediated by the ASE, ADF, ASG, and ASI neurons. A receptor can be expressed in AWA using the promoter region of the *odr-10* gene operably linked to the receptor coding region (Troemel ER, *et al*, *supra*). A receptor can be expressed in the ASE neuron using the promoter regions of the *gcy-5*, *gcy-6*, or *gcy-7* genes operably linked to the receptor coding region (Yu S, *et al*, *PNAS* 91:3384-3387 (1997)). Dauer formation is mediated by the ADF, ASI and ASJ neurons; a human 7TMR can be expressed in these neurons using the promoter region of the *gpa-10* gene operably linked to the receptor coding region (Jansen G, *et al*, *supra*). Thermotaxis is mediated by the AFD neuron; a human 7TMR can be expressed in this neuron using the promoter of the *gcy-8* gene operably linked to the receptor coding region (Yu S, *et al*, *supra*). Once a human 7TMR is expressed in the appropriate neuron to elicit a desired behavior, these behaviors can then be used to, for example, identify the substances that act as ligands to activate the human 7TMR, identify substances that antagonize the human 7TMR activation, determine the human 7TMR that is activated by a particular substance, or determine the potency and efficacy of activating substances.

In a particularly preferred embodiment of the invention, transgenic *C. elegans* are produced that express human 7TMRs in the sensory neurons that correlate with behavior, as discussed above. Such human 7TMRs are expressed in particular sensory neurons of in the transgenic *C. elegans* to select the desired behavior of the transgenic *C. elegans* upon human 7TMR activation. DNA constructs that direct expression of human 7TMRs will be composed of up to four different DNA segments: (1) a promoter segment; (2) an optional signal peptide encoding segment; (3) a receptor-encoding segment; and (4) and a 3' segment. These constructs are then introduced into nematodes by methods well known in the art to produce a stable strain that express the receptors in the appropriate sensory neurons.

The promoter DNA segment, importantly, confers the appropriate temporal and spatial characteristics to human 7TMR expression. In *C. elegans*, chemotactic behavior can

be defined in four broad categories, each mediated by a subset of sensory neurons.

Bargmann, *supra*. Genes that are expressed in particular sets of sensory neurons are known. Using recombinant DNA technique, the promoters or DNA segments that confer temporal/spatial specificity to these genes can be fused to a heterologous gene. Sambrook, *et al.*, 1. *Plasmid Vectors*. In Sambrook J, Fritsch EF, and Maniatis, T (eds), MOLECULAR CLONING: A LABORATORY MANUAL, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York. (1989).

For example, reporter gene expression in the ADL and ASH neurons, both of which mediate avoidance behavior when exposed to water-soluble substances, can be driven by the *gpa-11* promoter. Jansen, *et al.*, *Nature Genetics* 21: 414-419 (1999). Any such promoter or DNA regulatory element is suitable, provided it directs gene expression specifically, and solely, in sensory neurons that mediate a particular behavior. The promoter segment should include all necessary DNA to provide transcriptional initiation, 5' mRNA processing, and translational initiation at a methionine start codon.

In order to bind exogenously applied substances, human 7TMRs are, preferably, localized to the cell surface membrane of the expressing cells. Proteins can be directed to the cell surface by virtue of a signal peptide sequence that is found at the amino-terminal portion of the protein. Heijne, *Nucleic Acids Research* 14: 4683-4690 (1986). This signal peptide sequence can be introduced in one of two ways. First, the DNA segment encoding the receptor itself may already include a functional signal peptide sequence. Thus, directing the human 7TMR to the cell surface, where it can respond to extracellular molecules, can be accomplished by maintaining the peptide signal sequence of an inserted receptor protein coding sequence. Alternatively, a separate DNA segment that encodes a peptide signal sequence can be introduced between the promoter and receptor-encoding DNA.

Ideally, the receptor-encoding segment of DNA will include the nucleotides necessary to encode the complete human 7TMR peptide, from the start methionine codon to the translational termination, or stop, codon. The DNA segment should be operably linked to the promoter DNA segment such that translational initiation will begin at the start methionine codon of the receptor-encoding DNA segment.

A 3' DNA segment is necessary to provide the appropriate transcriptional termination and 3' mRNA processing signals. Although the 3' regions of many genes may be sufficient for this purpose, the standard DNA segment used for expression in *C. elegans* is composed of a 3' portion of the *unc-54* gene. Fire, *et al.*, *Gene* 93:189-198 (1990). See

Example 1, which depicts how to create a DNA construct that is capable of driving expression of a non-nematode cell surface receptor in the ADL and ASH neurons of *C. elegans* that mediate water-soluble chemorepulsion. Other vectors have been described that can drive expression in the AWA and AWB neurons (Troemel ER, *et al*, *Cell* 91:161-169 (1997)).

Plasmid constructs can be introduced into *C. elegans* using described transformation methods to generate animals transgenic for the plasmid constructs. Mello, *et al.*, *EMBO J.* 10: 3959-3970 (1991). Preferably, the plasmid used for this method is a linear plasmid. An important aspect of transformation in *C. elegans* is that plasmid constructs are easily cotransformed. Mello, *et al.*, *supra*. Therefore, transgenic strains of *C. elegans* can be generated that express more than one human 7TMR. In addition, dominant cotransformation markers can be used to score for transgenic *C. elegans*. In a particularly preferred embodiment of the invention, a cotransformation marker that does not affect the ability of the animals to exhibit chemotactic behavior is used. This cotransformation marker can be, for instance, a DNA fragment that complements a mutant phenotype. For example, mutations in the *C. elegans unc-119* gene cause a severe movement defect. Introduction of a segment of DNA containing the wildtype copy of the *unc-119* gene will complement the movement defect (Maduro, *Genetics* 141: 977-88 (1995)), causing the transgenic animals to be wildtype in movement. Cotransforming *unc-119* animals with the *unc-119* wildtype DNA and the sensory neuron expression plasmids will allow the presence of the transgene to be scored by rescue of the *unc-119* mutant phenotype. Alternatively, a reporter gene can be used that can be scored in a living animal, but does not affect the movement of the animal. For example, green fluorescent protein (herein referred to as "GFP") is a widely used reporter molecule in living systems. Ellenberg, *Trends Cell Biol.* 9(2): 52-56 (1999); Chalfie, *et al.*, *Science* 263(5148): 802-05 (1994). Co-injection of a GFP-expressing DNA construct with the sensory neuron expression plasmids would allow the scoring for the presence of the transgene by examination of animals for GFP fluorescence. Ellenberg, *et al.*, *supra*.

Therefore, in another aspect, this method of producing transgenic *C. elegans* further comprises the step of:

(c) introducing a DNA construct that expresses a reporter gene with a nematode promotor.

Preferably, this reporter gene encodes a fluorescent protein, such as a green fluorescent protein (GFP).

Expression and activation of a particular human 7TMR may not be sufficient to produce a behavioral response from a transgenic *C. elegans*. This may happen if the receptor requires accessory proteins to activate downstream signaling pathways. For example, many human 7TMRs activate downstream pathways through the action of heterotrimeric G proteins. Stadel, *et al.*, *supra*. Although *C. elegans* contains 20 G $\alpha$ , 2 G $\beta$ , and 2G $\gamma$  G proteins (Jansen, *et al.*, *supra*.), efficient activation of signaling pathways and subsequent behavioral response of the animal may require the coexpression of accessory proteins, such as G proteins. In a preferred embodiment of the invention, such accessory proteins are human G proteins. This can be accomplished using the same DNA constructs used to express the human 7TMR, replacing the receptor-encoding DNA segment with a DNA segment encoding the accessory protein. The constructs encoding the receptor and the accessory protein can then be cotransformed into nematodes.

A preferred embodiment of the invention provides a method for producing a transgenic *C. elegans* that expresses a human 7TMR in the sensory neurons that correlate with behavior, said method comprising the steps of:

- (a) producing a first transgene by operably linking a gene expression construct that encodes a human 7TMR to a sensory neuron promotor;
- (b) producing a second transgene by comprising an accessory protein operably linked to a promotor; and
- (c) introducing said first and second transgenes into said *C. elegans*, such that said *C. elegans* coexpresses both said accessory protein and said human 7TMR in the sensory neurons that correlate with behavior.

In a particularly preferred embodiment of this method, the accessory protein is a human G protein.

A preferred embodiment of the instant invention encompasses the transgenic *C. elegans* that are produced by the methods described above.

#### **Assaying Behavior in Transgenic *C. elegans***

Test populations of transgenic *C. elegans* that express human 7TMRs in their sensory neurons will be able to sense and react to molecules or substances that modify receptor activity. Depending on the sensory neuron(s) that the human 7TMR(s) are expressed in, the reaction may be movement toward, or movement away from, a source of the substance; thermotaxis behavior; or commitment to the dauer pathway. A large variety of assays can be configured based on these behaviors. We first describe ways of presenting

test substances to the animals, and then describe ways of measuring the behavior of the animals.

In one embodiment, the instant invention provides a method for identifying at least one ligand of at least one human 7TMR, said method comprising the steps of:

- 5 (a) providing at least one *C. elegans* that expresses at least one human 7TMR in the sensory neurons that correlate with behavior;
- (b) contacting said at least one *C. elegans* with at least two different concentrations of at least one test substance; and
- (c) detecting modulation of behavior of said at least one *C. elegans* in response to
- 10 said at least one test substance.

In a preferred embodiment of the invention, substances are presented to test populations of transgenic *C. elegans* in such a way that allows chemosensory behavior. In such chemotaxis assays, preferably, the animals are placed on a substrate that allows free movement, such as a solid or a semi-solid substrate. For example, a typical culture medium

15 for *C. elegans* is composed of a buffered solidified agar substrate placed in a container such as a petri plate. Sulston WB, *supra*. The substrate medium can be buffer or it can be growth medium, such as agar. In order to observe chemotactic behavior, it is preferable that a test substance be present in at least two different concentrations at spatially distinct regions of the substrate. A number of standard chemotactic assays have been described,

20 including assays that use point sources, gradients, and simple two concentration presentation of test substances. See Bargmann, *supra*; Troemel, *et al.*, *supra*; Jansen, *et al.*, *supra*. All of these assays have the following two characteristics: (1) a substrate medium allows the animals to locomote efficiently between, and show a preference for, different regions of the substrate; and (2) the presence of a test substance at two concentrations or

25 more in spatially distinct regions of the substrate medium.

Test substances can be presented in a dispersed format, or a point source format. An example of a dispersed format presentation method is the water-soluble chemorepulsive assay that is performed in divided petri dishes. Jansen, *et al.*, *supra*. In this format, a test substance is added to the agar poured in one half of a divided petri dish. In the other half of

30 the petri dish, agar without the substance is poured. A small drop of isoamyl alcohol, which is a strong diffusable chemoattractant to *C. elegans*, is placed on the agar half that contains the test substance and a population of test animals is placed on the opposite half. Other known chemoattractants can be used, such as biotin, lysine, histidine, cysteine, and serotonin in place of the isoamyl alcohol. Bargmann, *et al.*, *supra*. After a suitable time

period, typically 90 minutes, the dispersion of animals on each half of the plate is quantitated by one of the methods discussed above. The fraction of animals that remain on the non-substance containing agar half of the petri plate is a reflection of how well the test substance can "repel" the animals. In assays with strong chemorepellants, such as copper sulfate, sodium dodecyl sulfate (SDS), D-tryptophan, heptanol, octanol, nonanol, nonanone, benzaldehyde, 2, 4, 5-trimethylthiazole, and ethyl heptanoate, more than 90% of the animals typically remain on the half of the agar plate without the chemorepellant. Jansen, *et al, supra.*; Bargmann, *et al., supra.* The chemorepulsive assay could be used in conjunction with the pCEASH2.1 vector described in example 1, which directs expression in sensory neurons that mediate water-soluble chemorepulsion.

In a particularly preferred embodiment, the instant invention provides a method for identifying at least one ligand of at least one human 7TMR, said method comprising the steps of:

- (a) providing at least one *C. elegans* that expresses at least one human 7TMR in the sensory neurons that correlate with water-soluble chemorepulsive behavior;
- (b) placing a medium into a first portion of a receptacle, wherein said receptacle is divided into at least two portions;
- (c) adding at least two different concentrations of at least one chemorepulsant substance to the first portion of said receptacle;
- (d) adding said at least one *C. elegans* to said second portion of said receptacle; and
- (e) detecting, after a suitable time period, the behavioral response of said at least one *C. elegans* to said at least one chemorepulsant substance.

In yet another particularly preferred embodiment, the instant invention provides a method for identifying at least one ligand of at least one human 7TMR, said method comprising the steps of:

- (a) providing at least one *C. elegans* that expresses at least one human 7TMR in the sensory neurons that mediate water-soluble chemoattractive behavior;
- (b) placing a medium into a first portion of a receptacle, wherein said receptacle is divided into at least two portions;
- (c) adding at least two different concentrations of at least one chemoattractant substance to the first portion of said receptacle;
- (d) adding said at least one *C. elegans* to said second portion of said receptacle; and

(e) detecting, after a suitable time period, the behavioral response of said at least one *C. elegans* to said at least one test substance.

Point sources of substances could also be used to test chemorepulsion and/or chemoattraction. The square plate assay is an example of a point source assay used to measure both chemoattraction and chemorepulsion. *See* Troemel, *et al.*, *supra*. Two spots of test substance are placed on the surface at one end of a square agar plate. Two spots of a control substance are placed at the other end, and a test population of *C. elegans* is placed in the center of the plate. After a suitable time period, the distribution of animals on the plate is quantitated. If the test substance is chemoattractive, the animals will tend to move toward that end of the plate. If the test substance is chemorepulsive, the animals will move away from that end of the plate. Although a square plate facilitates quantitation of the animals, the shape is not important for behavior. Point sources of substances could be used with round plates, for example. Like the chemorepulsive assay described previously, this assay could be used in conjunction with receptor expression from pCEASH2.1. In addition, this assay could be used with other vectors that drive expression in neurons mediating water-soluble chemoattractive behavior, or chemotactic behavior toward volatile compounds.

In addition to previously described assays, we propose three novel variations that are particularly preferred embodiments of the invention. First, presentation of substances as point sources can be used to assay large numbers of test substances by creating an array of spots of at least one test substance on a substrate medium. *See* Figure 2. A number of different test substances can be placed in an array of spots on a plate. A test population of transgenic *C. elegans* created, as described above, can then be introduced and assayed to determine their chemotactic preference for the test substances. Spots of test substances that the *C. elegans* preferentially avoid, or are attracted to, would then be candidates for substances that modify the activity of human 7TMRs that are expressed in the sensory neurons of the test population of *C. elegans*. In a particularly preferred embodiment of the instant invention, the test substances can easily diffuse in the substrate medium. Presenting test substances as an array of point sources confers the advantage of being able to rapidly assay a large number of different point sources of substance for their effect on chemotactic behavior.

Example 2, detailed below, demonstrates a chemotactic assay using a simple array of substances. In this experiment, four different substances were placed in spot on a agar substrate. One substance, isoamyl alcohol, is a known strong chemoattractant to wildtype



*C. elegans*. Bargmann, *supra*. A test population of wildtype *C. elegans* was then placed at the center of the petri dish, equidistant from each of the substances. The position of the animals was photographed at the initial time they were placed on the substrate and at a later time. As can be seen, at the later time point, the animals preferentially moved to the

5 location of the strong chemoattractant. In a particularly preferred embodiment of our invention, the test population can be transgenic for a 7TMR gene, and the test substances can be molecules that potentially modify the activity of the receptor, as discussed above.

In yet another particularly preferred embodiment of the invention, microorganismal colonies, such as bacteria, viruses or yeast, are used as the point sources of test substances.

10 A large number of techniques exist to create "libraries" of substances expressed by microorganisms. Cesareni, *et al.*, *Comb Chem High Throughput Screen* 2(1): 1-17 (1999); Georgiou, *et al.*, *Nat Biotechnol.* 15(1): 29-34 (1997); Sambrook, *et al.*, 12. *Screening Expression Libraries with Antibodies and Oligonucleotides*. In Sambrook J, Fritsch EF, and Maniatis, T (eds), *MOLECULAR CLONING: A LABORATORY MANUAL*, Cold Spring

15 Harbor Laboratory Press, Cold Spring Harbor, New York. (1989). These libraries can be used in our inventive methods to identify specific test substances that modify activity of the human 7TMRs expressed in the test population of *C. elegans*. For example, microorganismal colonies, each of which secrete a different substance, can be placed on the substrate medium. A test population of *C. elegans* can then be introduced onto the

20 substrate. The behavioral preference of the animals to these colonies would indicate potential activity of the substance secreted by the colony on the human 7TMR(s) expressed in the test population.

Yet another particularly preferred embodiment of the invention is a chemotaxis assay that places a test population of *C. elegans* on a biomolecular separation matrix. Such

25 a biomolecular separation matrix could be in agarose or polyacrylamide, and the animals could then be placed on the matrix. By monitoring the behavior of the animals, any preference for certain areas of the matrix could be determined. For example, a tissue that might contain a ligand for a human 7TMR could be fractionated by molecular mass in a matrix-based separation system. The chemotactic behavior of animals that express the

30 human 7TMR in their sensory neurons could then indicate in what region of the matrix contained the ligand of the human 7TMR, and therefore, what mass the ligand was.

There are a large variety of separation techniques for biological samples, including those for small peptides, proteins, DNA, and other biological components. Many of these techniques use a matrix substrate that is amenable to a chemotaxis assay using test

populations of *C. elegans*. For example, nucleic acids and proteins can be separated in a simple agarose substrate. Sambrook, *et al.*, 6. *Gel Electrophoresis of DNA*, In Sambrook J, Fritsch EF, and Maniatis, T (eds), MOLECULAR CLONING: A LABORATORY MANUAL, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York. (1989);

5 Sambrook, *et al.*, 18 *Detection and Analysis of Proteins Expressed from Cloned Genes*. In Sambrook J, Fritsch EF, and Maniatis, T (eds), MOLECULAR CLONING: A LABORATORY MANUAL, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York (1989). Once the separation is complete, a test population of *C. elegans* can be introduced onto the separation matrix. Behavioral preference for, or avoidance of, different

10 regions of the matrix would indicate the presence of a substance at that location that modifies the activity of human 7TMR(s) expressed in the chemosensory neurons. The measurement of chemotactic behavior involves determining the spatial distribution of a test population of transgenic *C. elegans* of the instant invention at an initial time point and at a later timepoint when the test population is exposed to at least two spatially distinct

15 concentrations of a substance. Methods for assaying the distribution of *C. elegans* have already been described. See Bargmann, *supra*; Troemel, *et al.*, *Cell* 91:161-169 (1997); Jansen, *et al.*, *supra*. In addition to these methods, we describe two embodiments of the instant invention that determine the spatial distribution of animals in the application of the inventive screening methods disclosed herein.

20 The animals may be detected by direct observation. For example in the water-soluble chemorepulsive assay, described above, an initial population of animals is placed at a defined location. After a suitable time period, the position of the animals is determined by direct observation of the agar surface using a suitable optical instrument. A chemotactic index can then be defined as the fraction of animals on the half of the petri plate containing

25 the test substance (number of animals on test substance/total number of animals). The chemotactic index for the test substance can then be compared to the indices determined for compounds for which chemotactic behavior has been defined.

An alternative to direct observation is mechanical detection of the animals. For instance, such detection could involve the determination of optical density across the test

30 surface by a machine. The animals would be detected by changes in density at the location where an animal was located. Alternatively, if the animals are expressing a reporter gene that can be detected in living animals (*i.e.*, GFP), a machine could monitor the position of the animals using a suitable reporter gene detection protocol.

In a particularly preferred embodiment of the invention, determining spatial distribution of the transgenic *C. elegans* involves assaying the feeding behavior of the animals. Soil nematodes, such as *C. elegans*, typically feed on bacteria, and it is possible to grow *C. elegans* on a variety of bacterial species. Venette, *et al.*, *Soil Biology and Biochemistry* 30: 949-960 (1998). In a preferred embodiment of the invention, such bacteria are selected from the group consisting of: *Acinetobacter calcoaceticus*, *Bacillus cereus*, *Bacillus sp.*, *Enterobacter amnigenus*, *Enterobacter cloacae*, *Escherichia coli*, *Flavobacterium sp.*, *Proteus vulgaris*, *Pseudomonas aeruginosa*, *Pseudomonas cepacia*, *Pseudomonas maltophilia*, *Pseudomonas putida*, *Pseudomonas sp.*, *Serratia liquefaciens*, *Serratia marcescens*, *Zymobacillus macerans*. In the laboratory, strains of nematodes, including *C. elegans*, are customarily grown on uniform "lawns" of *E. coli* bacteria placed on an agar surface. Epstein, *et al.*, *CAENORHABDITIS ELEGANS: MODERN BIOLOGICAL ANALYSIS OF AN ORGANISM*, Academic Press, San Diego, CA. (1995).

As the *C. elegans* grow and reproduce, the *E. coli* lawn is diminished until no bacteria remain. Therefore, the reduction of the bacterial lawn is a measurement of an actively feeding population of *C. elegans*. The rate of decrease of the lawn is proportional to the number of feeding animals present. Example 3, below, demonstrates this assay. The consumption of bacteria can be used as an indirect indicator of the distribution of animals. Substances can be spotted onto a substrate in an array, and a uniform lawn of bacteria placed on the substrate, overlaying the array. A test population of *C. elegans* can be introduced and incubated for a suitable time period to allow consumption of the lawn. If the animals are attracted to a substance, then they will move to that region of the substrate and the bacterial lawn will be diminished at that spot in comparison to the surrounding lawn. Conversely, if a substance repels the animals, then the bacterial lawn at that spot will not be consumed. Thus, bacterial consumption can be used to assay the growth and location of a population of nematodes. If a GFP-expressing bacterial strain is used (Chalfie, *et al.*, *supra*) as was done in the experiment of Example 3, the variations in bacterial density can be easily detected, for example, by photographing fluorescence intensity.

In another particularly preferred embodiment of the invention, chemotactic behavior of the transgenic *C. elegans* is assayed by combining the ability of *C. elegans* to consume bacteria with the ability to generate recombinant libraries of substances in bacteria, as discussed above. For instance, a library of secreted substances can be engineered through recombinant DNA methods in bacteria. Georgiou, *et al.*, *supra*. The library can be plated out as individual colonies on an agar substrate, with each colony

secreting a different substance. A test population of *C. elegans* could then be introduced onto the substrate. With no chemotactic preference shown by the animals, the colonies can be consumed at the same rate. However, if one or more of the colonies expressed a substance that was chemorepulsive to the animals, then that colony or colonies can be consumed at a slower rate. These "resistant" colonies could then be recovered and the substance identified through analysis of the bacterial clone.

Thus, in a particularly preferred embodiment, the instant invention provides a method for identifying at least one ligand of a human 7TMR said method comprising the steps of:

- (a) providing at least one *C. elegans* that expresses a human 7TMR in the sensory neurons that correlate with behavior;
- (b) at least two different concentrations of at least one test substance onto a substrate surface that contains growth medium;
- (c) placing a uniform lawn of bacteria onto the surface of said growth medium;
- (d) contacting said at least one *C. elegans* with the said uniform lawn of bacteria; and
- (e) detecting, after a suitable time period, a decrease in the density of said uniform lawn of bacteria.

The dauer pathway is an alternative life cycle stage common to many nematode species. Riddle, *et al.*, *supra*. The dauer pathway is normally triggered by environmental stresses such as starvation, temperature extremes, or overcrowding. Genetically, the dauer pathway has been most intensively studied in *C. elegans*. The response to overcrowding in *C. elegans* is mediated by a substance known as dauer pheromone, which is secreted by the animals. When dauer pheromone becomes sufficiently concentrated, it triggers commitment to the dauer alternative life cycle stage. Riddle, *et al.*, *supra*. Three sensory neurons have role in the appropriate response to dauer pheromone: ADF, ASI, and ASJ neurons. Bargmann, *supra*. In addition to pheromone, activation of a transgenically expressed human 7TMR in these neurons of *C. elegans* would be expected to lead to commitment to the dauer pathway. Unlike chemotaxis assays, measuring dauer formation does not require presentation of substances on a substrate that allows movement of *C. elegans*, nor does it require that the substance be present at two spatially distinct concentrations. Preferably, the test population of *C. elegans* are exposed to the substance during the portion of the life cycle at which commitment to the dauer pathway is made. Riddle, *et al.*, *supra*. Measurement of dauer formation has been previously described. *Id.*

Thermotaxis chemosensory behavior can be used to assay the function of human 7TMRs that might be affected by temperature. A test population of *C. elegans* can be generated as described above that express a human 7TMR in the AFD neuron. The AFD neuron mediates thermotaxis, and expression of a thermoreactive human 7TMR in this

5 neuron can modify the thermotactic behavior of the test population of *C. elegans*. The test population of animals can then be placed on a solid substrate that is maintained in a thermal gradient, as described (Bargmann, *supra*). The preference of the test population for certain thermal isoclines on the substrate can be used as a measure of human 7TMR activation.

In a particularly preferred embodiment of the invention, the chemosensory behavior

10 of *C. elegans* can be used to identify substances that activate human 7TMRs. As outlined above, human 7TMRs can be expressed in *C. elegans*, and the *C. elegans* can then be exposed to an array of substances. By comparing the chemosensory behavior seen in test populations of wildtype animals (non-transgenic) and transgenic animals expressing a particular human 7TMR, substances can be identified that alter the behavior of only the

15 transgenic *C. elegans*. These substances would then be candidates for interacting with and/or modifying the activity of the transgenically-expressed human 7TMR. Thus, a particularly preferred embodiment of the invention encompasses screening for substances that can activate human 7TMRs that have no known ligand, which are referred to in the art as "orphan receptors". This will be especially important in light of determination of the

20 human genome sequence. Orphan 7TMRs can be expressed in *C. elegans* and, as outlined above, an array of test substances could be exposed to the animals. The behavior of the animals will then indicate the presence of an activating substance or ligand.

In yet another preferred embodiment of the invention, the test substances are present in beads, and the medium is solid or semi-solid.

25 Thus, a particularly preferred embodiment of the instant invention provides a method for identifying at least one ligand of at least one human 7TMR, said method comprising the steps of:

- (a) providing at least one *C. elegans* that expresses a human 7TMR in the sensory neurons that correlate with behavior;
- 30 (b) placing a medium in a receptacle;
- (c) placing at least two different concentrations of at least one test substance in said receptacle;
- (d) adding said at least one *C. elegans* to said receptacle; and

(e) detecting, after a suitable time period, the behavioral response of said at least one *C. elegans* over the surface of said medium.

Using this approach, the instant invention would allow the inexpensive testing of a large number of substances. For example, once a transgenic strain of *C. elegans* is generated, large populations of transgenic *C. elegans* can be grown for chemotactic assays. Many substances can be tested by spotting them in a large array on the substrate medium, as discussed above. In addition, a large variety of substances created by recombinant DNA techniques in microorganisms can be assayed at once. Thus, our invention can be used to identify rare or novel substances that act on certain human 7TMRs. This ability will have special utility when analyzing the human 7TMRs identified only through analysis of DNA sequence.

In a particularly preferred embodiment of the invention, screening methods with the transgenic *C. elegans* can be used to evaluate the potency and specificity of substances on human 7TMR activity. A strain of *C. elegans* can be generated that expresses a particular human 7TMR upon which a substance is known to act. Structurally related substances could then be tested and compared in a chemotactic assay for their efficacy in eliciting a chemotactic response from the human 7TMR-expressing strain. Because chemotactic response is a result of receptor activation, the response can be used as a gauge for how well substances can activate the human 7TMR. For instance, structurally related compounds can be tested to determine which chemical differences are important in activating the human 7TMR and eliciting a chemosensory response. Such information is important in determining the structure/activity relationship of different substances, and can lead to an understanding of what part of the chemical structure is important for receptor activity.

For example, a group of 12 receptors might be known, one of which interacts with a substance. Four transgenic strains of *C. elegans* can be generated that express three human 7TMRs each. These strains can be assayed for chemosensory response to the substance. The three human 7TMRs in the responding strain can then be expressed in individual strains to identify the responding human 7TMR.

In yet another preferred embodiment, the instant invention can be used to identify a human 7TMR activated by a particular substance by generating transgenic strains of *C. elegans* that express more than human 7TMR, as discussed above. For example, a large group of human 7TMRs may be identified, one of which is activated by a particular substance. Transgenic strains of *C. elegans* can be generated that express overlapping subsets of the human 7TMR group. By testing the chemosensory behavior of these strains

toward the substance, the subset of human 7TMRs common to the strains that respond to the substance can be identified. These human 7TMRs can be further subdivided and transgenic strains of *C. elegans* made to identify the particular human 7TMR that responds to the substance.

5 In a particularly preferred embodiment, the instant invention provides a method for evaluating the potency of human 7TMR activation by a known ligand, said method comprising the steps of:

- (a) providing at least one *C. elegans* that expresses at least human 7TMR, in the sensory neurons that correlate with behavior, wherein said human 7TMR has a known  
10 ligand;
- (b) contacting said at least one *C. elegans* with said ligand and at least one structurally related compound; and
- (c) detecting the behavioral response of said at least one *C. elegans* to said at least one structurally related compound; and
- 15 (d) comparing the behavioral response of said at least one *C. elegans* to said ligand to the behavioral response of said at least one *C. elegans* to said at least one structurally related compound.

In yet another particularly preferred embodiment of the invention, screening methods can be used to identify an antagonizing substance that blocks the effect of a  
20 substance on human 7TMR. For example, if a ligand-receptor pair is already known, our invention can be used to screen for or evaluate the ability of substances to antagonize receptor activation by the ligand. The chemosensory response of animals to the ligand in the presence and absence of substances can be assayed. If a substance antagonizes receptor activation by the ligand, then the chemotactic response of the animals to the ligand will be  
25 abolished in the presence of antagonist. If the test substance agonizes ligand activation of the receptor, then the chemotactic response should be enhanced in the presence of the substance.

This particularly preferred embodiment of the invention provides a method for identifying at least one test substance that modulates the activation of at least one human  
30 7TMR by a known ligand, said method comprising the steps of:

- (a) providing at least one *C. elegans* that expresses a human 7TMR in the sensory neurons that correlate with behavior, wherein said human 7TMR has a known ligand;
- (b) contacting said at least one *C. elegans* with at least one test substance in the presence of said ligand;

(c) measuring the alteration of behavior of said at least one *C. elegans* in response to said at least one test substance; and

(d) determining whether said at least one test compound binds to and activates or inhibits the activation of said human 7MR by said ligand.

5 In addition to chemotactic behavior, many other *C. elegans* behaviors are regulated by a variety of neurons within the nervous system. We propose that these behaviors can be modified by expression of a human 7TMR within these neurons in the nervous system, and that the behavior modification can be applied to the identification and characterization of substances that activate the human 7TMR, as outlined in the Field of the Invention.

10 Large-scale functional evaluation of the *C. elegans* genome has defined a number of standard phenotypes that are easily scored by trained *C. elegans* biologists: exploded (Exp), dumpy (Dpy), long body (Lon), hyperactive movement (Hpr), paralyzed (Prl), molt defect (Mlt), sterile (Ste), sick (Sck), body morphology defect (Bmd), vulvaless (Vul), slow growth (Gro), egg laying defect (Egl), larval arrest (Lva), larval lethal (Let), protruding  
 15 vulva (Pvl), multiple vulva (Muv), sterile progeny (Stp), small (Sma), clear (Clr), blistered (Bli), high incidence of male progeny (Him), roller (Rol), larval lethal (Lvl), uncoordinated (Unc), embryonic lethal (Emb) (Maduro, *et al.*, *Genetics* 141, 977-88 (1995); Piano, *et al.*, *Current Biology* 10, 1619-22 (2000); Gonczy, *et al.*, *Nature* 408, 331-6 (2000); Fraser *et al.*, *Nature* 408, 325-30 (2000)). Of these phenotypes, four can directly result from  
 20 modification of nervous system function by gene mutation or expression of modified proteins: Hpr, Prl, Egl, and Unc. For example, expression of activated G proteins in the nervous system can lead to an Egl phenotype; in addition, mutations in a number of human 7TMR signaling pathway proteins can lead to an Egl phenotype, or suppress the action of a second mutation that leads to the Egl (Wilkie, *Current Biology* 10, R853-6 (2000)).  
 25 Therefore, expression and activation of a human 7TMR could perturb the *C. elegans* nervous system and manifest or modify these phenotypes. The phenotypic read-out could then lead to evaluation of substances that would alter human 7TMR activation.

The Hpr, Prl, Egl and Unc phenotypes have been well documented in the literature and standard methods have been described for each of these phenotypes. Hyperactive  
 30 behavior (Mendel *et al.*, *Science* 267, 1652-5 (1995)) can be scored by examining the tracks of individual animals as they move across a solid agar medium. Hyperactive animals tend to have a "loopier" movement. The Paralyzed phenotype (Prl) is self-explanatory; the animals are capable of very little movement and tend to remain in one place for a long



period of time (Saifee, *et al.*, *Molecular Biology of the Cell* 9, 1235-52 (1998)). Animals exhibiting Egl, or egg laying, phenotype typically lay fewer eggs than normal. A severe Egl animal lays very few eggs and eventually is consumed by progeny that hatch within the animal (Wang *et al.*, *Current Topics in Developmental Biology* 51, 189-220 (2001)). There are a large variety of Uncoordinated (Unc) phenotypes (Sulston, WB (ed), THE NEMATODE *CAENORHABDITIS ELEGANS*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, (1988)). In general, uncoordinated behavior can be any deviation from the smooth sinusoidal movement and reaction to prodding exhibited by a normal animal. The movement defects seen in the Hpr, Prl and Unc phenotypes have obvious links to the animals' nervous systems. A correctly functioning neural network is required to produce smooth movements and responses to stimuli in the environment. The Egl phenotype can result from defects in a set of neurons called the HSNs, which innervate the egg-laying muscles. All of these phenotypes are well-characterized and can readily be identified by a trained *C. elegans* biologist (Sulston, WB (ed), THE NEMATODE *CAENORHABDITIS ELEGANS*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, (1988)).

Perturbation of the *C. elegans* nervous system by a human 7TMR requires expression of the receptor of interest within the animal's nervous system. Such expression can be accomplished by fusing a DNA promoter segment that directs expression throughout the nervous system, *i.e.*, in all neurons (pan-neuronal expression), to a DNA segment that encodes the human 7TMR. The promoter of the *unc-119* gene is a well-characterized example of a pan-neuronal promoter (Maduro, *et al.*, *supra*). Example 4 details the construction and use of a plasmid that will drive pan-neuronal expression of a human 7TMR using the *unc-119* promoter DNA segment. This plasmid can be introduced via standard methods to produce a transgenic strain of *C. elegans* that express a human 7TMR throughout the nervous system (Mello, *et al.*, *EMBO Journal* 10, 3959-70 (1991); Praitis *et al.*, *Genetics* 157, 1217-26 (2001)). The resulting strains can then be examined for phenotypes, as detailed above. *C. elegans* Expression of a human 7TMR in the *C. elegans* nervous system can cause a phenotype-producing perturbation in at least three ways. First, the human 7TMR might be activated in one or many neurons by an endogenous ligand that is produced by the nematode. Activation of the receptor would lead to modification of neuronal function, thereby leading to an altered phenotypic state that can be assayed (see above). Second, the human 7TMR might be activated by an exogenously applied substance, again leading to modification of neuronal function and a resulting

phenotype. Third, although not activated, the human 7TMR might associate with accessory proteins, such as human G proteins, thereby modifying neuronal function and leading to a phenotype in the transgenic animal.

Phenotypes induced by any of the three mechanisms detailed above can be used to evaluate substances, as detailed in the Field of the Invention. For example, if a human 7TMR is activated by an endogenous ligand to produce a phenotype, that phenotype can be used to screen for and identify substances that would antagonize the receptor. Specifically, a strain of animals that express a certain receptor might exhibit a phenotype due to activation by an endogenous ligand. An exogenous substance that would block activation of the 7TMR would be able to suppress, or return to normal, the phenotype. To find such antagonistic substances, the strain expressing the human 7TMR could then be cultured in the presence of a variety of substances to see which would suppress the phenotype. These substances would then be candidates for molecules that would antagonize human 7TMR activation. Because the phenotypic assay does not involve chemotaxis behavior as described previously, the substances would not have to be spatially distributed and could be uniformly distributed in the medium.

In addition to identifying antagonistic substances, molecular genetic techniques could be used to identify a surrogate, ligand present in a human 7TMR-expressing strain that exhibits a phenotype. For example, a human 7TMR-expressing strain could be subjected to mutagenic screens to identify mutations that would suppress the phenotype. Such mutations would be candidates for lesions that eliminate production of the endogenous ligand. The mutations could be genetically mapped and molecularly cloned. Such suppressor genetic screens are commonly used in the field of genetics in a variety of contexts (Dong, *et al.*, *Genes & Development* 14, 2003-14 (2000)). Mapping and cloning techniques for mutations are also well-known and characterized (Wicks *et al.*, *Nature Genetics* 28, 160-4 (2001)).

Human 7TMR-expressing strains of *C. elegans* could also be used to find substances that would activate, or agonize, the receptor. For example, a strain of nematodes expressing a human 7TMR might not exhibit a phenotype because no endogenous ligand produced by the receptor-expressing strain activates the receptor. However, an exogenously applied agonistic substance that activated the human 7TMR might cause a receptor-expressing strain of nematodes to exhibit a phenotype. Therefore, agonistic substances could be identified by culturing the human 7TMR-expressing strain in

the presence of a variety of substances to see if such exposure leads to a phenotype. Substances identified in this way would be candidates for molecules that bind to and activate the human 7TMR.

A human 7TMR-expressing strain that exhibits a phenotype upon exposure an agonistic substance could be used to evaluate the specificity and efficacy of different substances. For example, structurally related substances could be evaluated for their effectiveness by comparing how well they elicit a phenotype from the same human 7TMR-expressing strain. Likewise, a human 7TMR-expressing strain that exhibits a phenotype suppressable by an exogenously applied antagonist could be used to evaluate different antagonistic substances. Thus, human 7TMR-expressing strains would provide a means to evaluate the specificity and efficacy of substances that interact with the human 7TMRs.

In summary, the invention proposes that expression of human 7TMRs within the neurons of *C. elegans* can modify behaviors and create phenotypes. These behaviors and phenotypes are useful for the identification of agonist and antagonist substances that act on the human 7TMRs. In addition, the behaviors and phenotypes are useful for the evaluation of specificity and efficacy of substances that interact with human 7TMRs.

## Examples

### Example 1: DNA Construct for Production of Transgenic Nematodes that Express Cell

#### Surface Receptors:

Figure 1 diagrams a constructed plasmid, named pCEASH2.1, capable of driving expression in the ADL and ASH neurons of *C. elegans* that mediate water-soluble chemorepulsion. Basepairs 1-7 correspond to the unique Kpn I site in the multiple cloning sites (MCS) region of the plasmid pBSKS(+) (Stratagene; Genbank Accession Number X52331). Basepairs 7-2830 correspond to nucleotides 31002 to 33825 in the sequence of cosmid C16A11 (Genbank Accession Number AF077536), with the exception that nucleotides 33818-33825 have been altered to read "CCATGGCC". This 2824 basepair DNA segment corresponds to the promoter region of the *C. elegans gpa-11* gene, up to and including the start methionine codon. The alterations to the sequence result in the introduction of a NcoI site at the start codon of the *gpa-11* protein coding region. Basepairs 2831 to 2907 correspond to the MCS region of pBSKS(+), from the unique Xho I site to the unique Not I site. Basepairs 2908 to 4178 correspond to basepairs 30364 to 29100 in the sequence of cosmid F32A7 (Genbank Accession Number Z83107), except for the addition

of the sequence "AATT" at position 29553 in the F32A7 sequence (this change eliminates an EcoRI site). This DNA segment contains the last 15 codons of the *C. elegans* unc-54 protein coding region, followed by 1228 basepairs of DNA 3' to the unc-54 stop codon. This region provides transcriptional termination and mRNA processing signals. Jansen, *et al.*, *supra*. Nucleotides 4179 to 7036 correspond to the pBSKS(+) plasmid from the unique Sac I site in the polylinker up to, but not including, the unique Kpn I site. In summary, pCEASH2.1 contains the *C. elegans gpa-11* promoter sequence fused to the MCS of the pBSKS(+) plasmid, followed by the 3' region of the *C. elegans* unc-54 gene.

When introduced into *C. elegans*, the open reading frame located between the *gpa-11* initiation codon and the unc-54 stop codon is expressed in the ASH and ADL chemosensory neurons. A heterologous sequence can be introduced into this reading frame using the unique sites contained in the MCS of pCEASH2.1. If a human 7TMR protein coding sequence is engineered with an NcoI site at the start methionine, it can be fused to the *gpa-11* start methionine such that translational initiation will occur at the start methionine of the inserted sequence. In this way, the authentic peptide signal sequence of a receptor can be maintained as discussed above, while the appropriate DNA regulatory elements are introduced to direct expression of the human 7TMR protein in the ASH and ADL sensory neurons.

#### 20 Example 2: Chemotaxis Assay

Figure 2 depicts a simple demonstration of a chemotaxis assay using an array of substances. In panel A, images of the experiment are shown. In panel B, a schematic of the images are drawn, with the positions of the test animals marked with black dots. The substrate is composed of a 0.5 cm. thick solidified agar solution (Troemel, *et al.*, *supra.*) poured in a petri dish that is 10 cm. in diameter. 4 microliter samples of the substances labeled in the schematic were placed equidistant from the center of the agar surface. A test population of *C. elegans* animals was placed in the center of the agar surface. Images were taken at the initial time and two hours after placement of the test population. The position of the animals at the two times is diagrammed in Figure 2. After two hours, the animals have preferentially moved toward the spot of isoamyl alcohol, a strong chemoattractant. Panel B depicts this movement.

Example 3: Bacterial consumption of a culture of GFP-expressing bacteria by *C. elegans*

In this experiment, GFP-expressing bacteria are placed in two microtiter wells along with a population of *C. elegans*. Because the bacteria express GFP, the amount of bacteria can easily be assayed using a fluorescent plate reader (Cytofluor, Perspective Biosystems). In one well, the medium also contains ivermectin, a nematocidal compound that kills *C. elegans*. Initially, both wells show equivalent levels of bacteria. However, after 4 days of culture, the bacteria in the well lacking ivermectin has been reduced, while the ivermectin-containing well shows bacterial growth. Figure 3 depicts this experiment. Thus, bacterial consumption can be used to assay the growth and feeding of a population of nematodes. If a GFP-expressing bacterial strain is used (Chalfie, *et al.*, *supra*) as was done in the experiment shown in Figure 3, the variations in bacterial density can be easily visualized by photographing fluorescence intensity.

Example 4: DNA Construct for Production of Transgenic *C. elegans* that Express Human 7TMRs within the Nervous System:

Figure 4 diagrams a constructed plasmid, named pCEPN, capable of driving pan-neuronal expression in *C. elegans*. Basepairs 1-7 correspond to the unique Kpn I site in the multiple cloning sites (MCS) region of the plasmid pBSKS(+) (Stratagene; Genbank Accession Number X52331). Basepairs 7-2205 correspond to nucleotides 1318 to 3516 in the sequence of *C. elegans* cosmid M142 (Genbank Accession Number Z73428), with the exception that nucleotides 3509-3516 have been altered to read "CCATGGCC". This 2199 basepair DNA segment corresponds to the promoter region of the *C. elegans* unc-119 gene, up to and including the start methionine codon (Maduro, Pilgrim, Genetics 141, 977-88 (1995)). The alterations to the sequence result in the introduction of a NcoI site at the start codon of the unc-119 protein coding region. Basepairs 2206 to 2282 correspond to the MCS region of pBSKS(+), from the unique Xho I site to the unique Not I site. Basepairs 2283 to 3553 correspond to basepairs 30364 to 29100 in the sequence of cosmid F32A7 (Genbank Accession Number Z83107), except for the addition of the sequence "AATT" at position 29553 in the F32A7 sequence (this change eliminates an EcoRI site). This DNA segment contains the last 15 codons of the *C. elegans* unc-54 protein coding region, followed by 1228 basepairs of DNA 3' to the unc-54 stop codon. This region provides transcriptional termination and mRNA processing signals. Jansen, *et al.*, *supra*. Nucleotides 3554 to 6411 correspond to the pBSKS(+) plasmid from the unique Sac I site in the polylinker up to, but not including, the unique Kpn I site. In summary, pCEPN

contains the *C. elegans* unc-119 promoter sequence fused to the MCS of the pBSKS(+) plasmid, followed by the 3' region of the *C. elegans* unc-54 gene.

When introduced into *C. elegans*, the open reading frame located between the unc-119 initiation codon and the unc-54 stop codon is expressed pan-neuronally (Maduro, Pilgrim, *Genetics* 141: 977-88 (1995)). A heterologous sequence can be introduced into this reading frame using the unique sites contained in the MCS of pCEPN. If a human 7TMR protein coding sequence is engineered with an NcoI site at the start methionine, it can be fused to the unc-119 start methionine, such that translational initiation will occur at the start methionine of the inserted sequence. In this way, the authentic peptide signal sequence of a receptor can be maintained as discussed above, while the appropriate DNA regulatory elements are introduced to direct expression of the human 7TMR protein in all neurons of the transgenic animal.

#### Example 5: Egg-laying (Egl) Phenotype Produced by Expressing 7TMRs Pan-Neuronally in *C. elegans*

DNA segments encoding six 7TMRs were cloned into the pCEPN expression vector (see Example 4). The five human 7TMRs are: GPR18 (SwissProt accession# Q14330), GPR7 (SwissProt accession# P48145), HCEPR (SwissProt accession# O00143), AXOR35 (SwissProt accession# Q9GZQ0), Octoray (tremble accession# Q9BY21). Transgenic *C. elegans* strains were created with the cloned receptor constructs using standard methods (Mello, *et al.*, *EMBO Journal* 10, 3959-70 (1991)). 5-8 single adult transgenic animals were picked from each strain and allowed lay eggs for 40 hours. The adult animals were removed and the number of eggs laid was quantitated. Figure 5 illustrates the results obtained. The normal, wildtype, strain of *C. elegans*, N2, laid 109 eggs, plus or minus 16, within the 40 hour period. In contrast, most of the human 7TMR-expressing strains reduce the number of eggs laid. In two cases, this Egl phenotype was relatively mild, with a reduction of around 50% in the number of eggs laid (GPR18 - 47% wildtype, GPR7 - 58% wildtype). However, two receptor-expressing strains showed a reduction in eggs laid of greater than 66% (HCEPR - 27% wildtype, AXOR35 - 34% and Octoray - 17%). In particular, Octoray exhibits a penetrant Egl phenotype, with Octoray-expressing animals laying less than 17% the normal number. This example shows that expression of non-nematode human 7TMRs in *C. elegans* can cause a scorable phenotype.

All publications including, but not limited to, patents and patent applications, cited in this specification, are herein incorporated by reference as if each individual publication were specifically and individually indicated to be incorporated by reference herein as though fully set forth.

- 5           The above description fully discloses the invention, including preferred embodiments thereof. Modifications and improvements of the embodiments specifically disclosed herein are within the scope of the following claims. Without further elaboration, one skilled in the art can, using the preceding description, utilize the present invention to its fullest extent. Therefore, the examples provided herein are to be construed as merely
- 10   illustrative and are not a limitation of the scope of the present invention in any way. The embodiments of the invention in which an exclusive property or privilege is claimed are defined as follows.

11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000  
1001  
1002  
1003  
1004  
1005  
1006  
1007  
1008  
1009  
1010  
1011  
1012  
1013  
1014  
1015  
1016  
1017  
1018  
1019  
1020  
1021  
1022  
1023  
1024  
1025  
1026  
1027  
1028  
1029  
1030  
1031  
1032  
1033  
1034  
1035  
1036  
1037  
1038  
1039  
1040  
1041  
1042  
1043  
1044  
1045  
1046  
1047  
1048  
1049  
1050  
1051  
1052  
1053  
1054  
1055  
1056  
1057  
1058  
1059  
1060  
1061  
1062  
1063  
1064  
1065  
1066  
1067  
1068  
1069  
1070  
1071  
1072  
1073  
1074  
1075  
1076  
1077  
1078  
1079  
1080  
1081  
1082  
1083  
1084  
1085  
1086  
1087  
1088  
1089  
1090  
1091  
1092  
1093  
1094  
1095  
1096  
1097  
1098  
1099  
1100  
1101  
1102  
1103  
1104  
1105  
1106  
1107  
1108  
1109  
1110  
1111  
1112  
1113  
1114  
1115  
1116  
1117  
1118  
1119  
1120  
1121  
1122  
1123  
1124  
1125  
1126  
1127  
1128  
1129  
1130  
1131  
1132  
1133  
1134  
1135  
1136  
1137  
1138  
1139  
1140  
1141  
1142  
1143  
1144  
1145  
1146  
1147  
1148  
1149  
1150  
1151  
1152  
1153  
1154  
1155  
1156  
1157  
1158  
1159  
1160  
1161  
1162  
1163  
1164  
1165  
1166  
1167  
1168  
1169  
1170  
1171  
1172  
1173  
1174  
1175  
1176  
1177  
1178  
1179  
1180  
1181  
1182  
1183  
1184  
1185  
1186  
1187  
1188  
1189  
1190  
1191  
1192  
1193  
1194  
1195  
1196  
1197  
1198  
1199  
1200  
1201  
1202  
1203  
1204  
1205  
1206  
1207  
1208  
1209  
1210  
1211  
1212  
1213  
1214  
1215  
1216  
1217  
1218  
1219  
1220  
1221  
1222  
1223  
1224  
1225  
1226  
1227  
1228  
1229  
1230  
1231  
1232  
1233  
1234  
1235  
1236  
1237  
1238  
1239  
1240  
1241  
1242  
1243  
1244  
1245  
1246  
1247  
1248  
1249  
1250  
1251  
1252  
1253  
1254  
1255  
1256  
1257  
1258  
1259  
1260  
1261  
1262  
1263  
1264  
1265  
1266  
1267  
1268  
1269  
1270  
1271  
1272  
1273  
1274  
1275  
1276  
1277  
1278  
1279  
1280  
1281  
1282  
1283  
1284  
1285  
1286  
1287  
1288  
1289  
1290  
1291  
1292  
1293  
1294  
1295  
1296  
1297  
1298  
1299  
1300  
1301  
1302  
1303  
1304  
1305  
1306  
1307  
1308  
1309  
1310  
1311  
1312  
1313  
1314  
1315  
1316  
1317  
1318  
1319  
1320  
1321  
1322  
1323  
1324  
1325  
1326  
1327  
1328  
1329  
1330  
1331  
1332  
1333  
1334  
1335  
1336  
1337  
1338  
1339  
1340  
1341  
1342  
1343  
1344  
1345  
1346  
1347  
1348  
1349  
1350  
1351  
1352  
1353  
1354  
1355  
1356  
1357  
1358  
1359  
1360  
1361  
1362  
1363  
1364  
1365  
1366  
1367  
1368  
1369  
1370  
1371  
1372  
1373  
1374  
1375  
1376  
1377  
1378  
1379  
1380  
1381  
1382  
1383  
1384  
1385  
1386  
1387  
1388  
1389  
1390  
1391  
1392  
1393  
1394  
1395  
1396  
1397  
1398  
1399  
1400  
1401  
1402  
1403  
1404  
1405  
1406  
1407  
1408  
1409  
1410  
1411  
1412  
1413  
1414  
1415  
1416  
1417  
1418  
1419  
1420  
1421  
1422  
1423  
1424  
1425  
1426  
1427  
1428  
1429  
1430  
1431  
1432  
1433  
1434  
1435  
1436  
1437  
1438  
1439  
1440  
1441  
1442  
1443  
1444  
1445  
1446  
1447  
1448  
1449  
1450  
1451  
1452  
1453  
1454  
1455  
1456  
1457  
1458  
1459  
1460  
1461  
1462  
1463  
1464  
1465  
1466  
1467  
1468  
1469  
1470  
1471  
1472  
1473  
1474  
1475  
1476  
1477  
1478  
1479  
1480  
1481  
1482  
1483  
1484  
1485  
1486  
1487  
1488  
1489  
1490  
1491  
1492  
1493  
1494  
1495  
1496  
1497  
1498  
1499  
1500  
1501  
1502  
1503  
1504  
1505  
1506  
1507  
1508  
1509  
1510  
1511  
1512  
1513  
1514  
1515  
1516  
1517  
1518  
1519  
1520  
1521  
1522  
1523  
1524  
1525  
1526  
1527  
1528  
1529  
1530  
1531  
1532  
1533  
1534  
1535  
1536  
1537  
1538  
1539  
1540  
1541  
1542  
1543  
1544  
1545  
1546  
1547  
1548  
1549  
1550  
1551  
1552  
1553  
1554  
1555  
1556  
1557  
1558  
1559  
1560  
1561  
1562  
1563  
1564  
1565  
1566  
1567  
1568  
1569  
1570  
1571  
1572  
1573  
1574  
1575  
1576  
1577  
1578  
1579  
1580  
1581  
1582  
1583  
1584  
1585  
1586  
1587  
1588  
1589  
1590  
1591  
1592  
1593  
1594  
1595  
1596  
1597  
1598  
1599  
1600  
1601  
1602  
1603  
1604  
1605  
1606  
1607  
1608  
1609  
1610  
1611  
1612  
1613  
1614  
1615  
1616  
1617  
1618  
1619  
1620  
1621  
1622  
1623  
1624  
1625  
1626  
1627  
1628  
1629  
1630  
1631  
1632  
1633  
1634  
1635  
1636  
1637  
1638  
1639  
1640  
1641  
1642  
1643  
1644  
1645  
1646  
1647  
1648  
1649  
1650  
1651  
1652  
1653  
1654  
1655  
1656  
1657  
1658  
1659  
1660  
1661  
1662  
1663  
1664  
1665  
1666  
1667  
1668  
1669  
1670  
1671  
1672  
1673  
1674  
1675  
1676  
1677  
1678  
1679  
1680  
1681  
1682  
1683  
1684  
1685  
1686  
1687  
1688  
1689  
1690  
1691  
1692  
1693  
1694  
1695  
1696  
1697  
1698  
1699  
1700  
1701  
1702  
1703  
1704  
1705  
1706  
1707  
1708  
1709  
1710  
1711  
1712  
1713  
1714  
1715  
1716  
1717  
1718  
1719  
1720  
1721  
1722  
1723  
1724  
1725  
1726  
1727  
1728  
1729  
1730  
1731  
1732  
1733  
1734  
1735  
1736  
1737  
1738  
1739  
1740  
1741  
1742  
1743  
1744  
1745  
1746  
1747  
1748  
1749  
1750  
1751  
1752  
1753  
1754  
1755  
1756  
1757  
1758  
1759  
1760  
1761  
1762  
1763  
1764  
1765  
1766  
1767  
1768  
1769  
1770  
1771  
1772  
1773  
1774  
1775  
1776  
1777  
1778  
1779  
1780  
1781  
1782  
1783  
1784  
1785  
1786  
1787  
1788  
1789  
1790  
1791  
1792  
1793  
1794  
1795  
1796  
1797  
1798  
1799  
1800  
1801  
1802  
1803  
1804  
1805  
1806  
1807  
1808  
1809  
1810  
1811  
1812  
1813  
1814  
1815  
1816  
1817  
1818  
1819  
1820  
1821  
1822  
1823  
1824  
1825  
1826  
1827  
1828  
1829  
1830  
1831  
1832  
1833  
1834  
1835  
1836  
1837  
1838  
1839  
1840  
1841  
1842  
1843  
1844  
1845  
1846  
1847  
1848  
1849  
1850  
1851  
1852  
1853  
1854  
1855  
1856  
1857  
1858  
1859  
1860  
1861  
1862  
1863  
1864  
1865  
1866  
1867  
1868  
1869  
1870  
1871  
1872  
1873  
1874  
1875  
1876  
1877  
1878  
1879  
1880  
1881  
1882  
1883  
1884  
1885  
1886  
1887  
1888  
1889  
1890  
1891  
1892  
1893  
1894  
1895  
1896  
1897  
1898  
1899  
1900  
1901  
1902  
1903  
1904  
1905  
1906  
1907  
1908  
1909  
1910  
1911  
1912  
1913  
1914  
1915  
1916  
1917  
1918  
1919  
1920  
1921  
1922  
1923  
1924  
1925  
1926  
1927  
1928  
1929  
1930  
1931  
1932  
1933  
1934  
1935  
1936  
1937  
1938  
1939  
1940  
1941  
1942  
1943  
1944  
1945  
1946  
1947  
1948  
1949  
1950  
1951  
1952  
1953  
1954  
1955  
1956  
1957  
1958  
1959  
1960  
1961  
1962  
1963  
1964  
1965  
1966  
1967  
1968  
1969  
1970  
1971  
1972  
1973  
1974  
1975  
1976  
1977  
1978  
1979  
1980  
1981  
1982  
1983  
1984  
1985  
1986  
1987  
1988  
1989  
1990  
1991  
1992  
1993  
1994  
1995  
1996  
1997  
1998  
1999  
2000  
2001  
2002  
2003  
2004  
2005  
2006  
2007  
2008  
2009  
2010  
2011  
2012  
2013  
2014  
2015  
2016  
2017  
2018  
2019  
2020  
2021  
2022  
2023  
2024  
2025  
2026  
2027  
2028  
2029  
2030  
2031  
2032  
2033  
2034  
2035  
2036  
2037  
2038  
2039  
2040  
2041  
2042  
2043  
2044  
2045  
2046  
2047  
2048  
2049  
2050  
2051  
2052  
2053  
2054  
2055  
2056  
2057  
2058  
2059  
2060  
2061  
2062  
2063  
2064  
2065  
2066  
2067  
2068  
2069  
2070  
2071  
2072  
2073  
2074  
2075  
2076  
2077  
2078  
2079  
2080  
2081  
2082  
2083  
2084  
2085  
2086  
2087  
2088  
2089  
2090  
2091  
2092  
2093  
2094  
2095  
2096  
2097  
2098  
2099  
2100  
2101  
2102  
2103  
2104  
2105  
2106  
2107  
2108  
2109  
2110  
2111  
2112  
2113  
2114  
2115  
2116  
2117  
2118  
2119  
2120  
2121  
2122  
2123  
2124  
2125  
2126  
2127  
2128  
2129  
2130  
2131  
2132  
2133  
2134  
2135  
2136  
2137  
2138  
2139  
2140  
2141  
2142  
2143  
2144  
2145  
2146  
2147  
2148  
2149  
2150  
2151  
2152  
2153  
2154  
2155  
2156  
2157  
2158  
2159  
2160  
2161  
2162  
2163  
2164  
2165  
2166  
2167  
2168  
2169  
2170  
2171  
2172  
2173  
2174  
2175  
2176  
2177  
2178  
2179  
2180  
2181  
2182  
2183  
2184  
2185  
2186  
2187  
2188  
2189  
2190  
2191  
2192  
2193  
2194  
2195  
2196  
2197  
2198  
2199  
2200  
2201  
2202  
2203  
2204  
2205  
2